

EXERCISE 1

Date issued: 17th April 2023

Date due: 25th April 2023

Homework Problem 1.1 (Miscellaneous on Convergence Rates)

11 Points

- (i) Explain why the definition of Q-quadratic convergence of a sequence requires the initial assumption that the sequence converges at all.
- (ii) Show that Q-quadratic convergence implies Q-superlinear convergence which implies Q-linear convergence which implies convergence.
- (iii) (a) Show that the notions of Q-linear, Q-superlinear and Q-quadratic convergence of a sequence imply their respective R-convergence counterparts.
(b) Give an example that shows that R-convergence of any kind of a sequence generally does not imply the corresponding Q-convergence.
- (iv) (a) Let $\|\cdot\|_a, \|\cdot\|_b: \mathbb{R}^n \rightarrow \mathbb{R}$ be equivalent norms. Show that Q-superlinear resp. Q-quadratic convergence of a sequence w.r.t. $\|\cdot\|_a$ implies Q-superlinear resp. Q-quadratic convergence w.r.t. $\|\cdot\|_b$.
(b) Give an example that shows that the corresponding statement can not hold for Q-linear convergence. Does it hold for R-linear convergence?

Homework Problem 1.2

(Visualizing and Interpreting Convergence Rates)

7 Points

- (i) For each of the following cases, give an example of a null sequence $(x^{(k)})$ in $(\mathbb{R}, |\cdot|)$ that
- (a) converges, but does not converge Q-linearly,
 - (b) converges Q-linearly, but does not converge Q-superlinearly,
 - (c) converges Q-superlinearly, but does not converge Q-quadratically,
 - (d) converges Q-quadratically, but does not converge with higher order.
- (ii) Explain what the Q-convergence rates of a sequence $x^k \rightarrow x^*$ will look like in a y -semi-logarithmic plot, i. e., when plotting the map $k \mapsto \ln |x^{(k)} - x^*|$.
- (iii) Plot the distance to the limit for the sequences from **task (i)** over the iterations in a standard and a y -semi-logarithmic plot. What do you observe?

Homework Problem 1.3 (Optimality Condition Gap)

9 Points

Consider the optimization problem

$$\text{Minimize } f(x) = (x_1 - x_2^2)(2x_1 - x_2^2) = 2x_1^2 - 3x_1x_2^2 + x_2^4 \quad \text{where } x \in \mathbb{R}^2.$$

- (i) Show that the necessary optimality conditions of first and second order are satisfied at $(0, 0)^T$.
- (ii) Show that $(0, 0)^T$ is a local minimizer for f along every straight line passing through $(0, 0)$.
- (iii) Show that $(0, 0)^T$ is not a local Minimizer of f on \mathbb{R}^2 .

Homework Problem 1.4 (First Order Conditions are Sufficient for Convex Functions)

2 Points

Let $f: \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function that is differentiable at $x \in \mathbb{R}^n$ with $f'(x) = 0$. Show that x is a global minimizer of f .

Please submit your solutions as a single pdf and an archive of programs via [moodle](#).